


Chemistry 12  
Review Sheet on Unit 2  
Chemical Equilibrium

1. What two things are equal at *equilibrium*?

\_\_\_\_\_ and \_\_\_\_\_

2. Consider the following *potential energy* diagram:




- a) Which reaction, forward or reverse, will be affected *more* by an increase in temperature? \_\_\_\_\_
- b) Write a thermochemical equation for the forward reaction using the numerical value for the heat.  
Answer \_\_\_\_\_
- c) When the temperature is first raised, which reaction will increase *most* in rate, forward or reverse? \_\_\_\_\_  
Explain why. \_\_\_\_\_
- d) If the rate of the forward reaction is *faster* than the reverse reaction for awhile, what will happen to the  $[A_2]$  and  $[B_2]$ ? \_\_\_\_\_


## Chemistry 12

## Unit 2 - Chemical Equilibrium

- e) If the  $[A_2]$  and  $[B_2]$  increases, what will happen to the rate of the *reverse reaction*?  
\_\_\_\_\_
- f) When the reverse reaction rate *catches up* to the forward reaction rate, the system is again at \_\_\_\_\_
- g) Since, for awhile, *the rate of the forward reaction was faster than the rate of the reverse reaction*, there would be an *increase* in the concentrations of \_\_\_\_\_ and a *decrease* in the concentration of \_\_\_\_\_ in the second equilibrium.
- h) We can summarize by saying that the equilibrium has shifted to the \_\_\_\_\_ as a result of increasing the temperature.
- i) Draw a graph showing the **rates** of the *forward* and *reverse* reactions vs. time summarizing what happens in 2(c) to 2(f).



3. Consider the reaction:



- a) If one mole of  $A_2$  and one mole of  $B_2$  are placed in a 1.0 L container, an equilibrium is established in which  $[A_2]$  and  $[B_2] = 0.40 \text{ M}$  and  $[C] = 1.2 \text{ M}$ . If 2.0 moles of C are placed into another 1.0 L container at the same temperature, what will the final concentrations of all the species be? (*HINT: This is not a calculation. It deals with how equilibrium can be approached from the left or from the right.* )

$[A_2] =$  \_\_\_\_\_  $[B_2] =$  \_\_\_\_\_  $[C] =$  \_\_\_\_\_

## Chemistry 12

## Unit 2 - Chemical Equilibrium

- b) Sketch **two** graphs showing each of the activities performed in 3a. The graphs are concentration vs. time.

(Starting with 1 mole  $A_2$  and 1 mole of  $B_2$ )



(Starting with 2 moles of  $C$ )




4. Give **four** characteristics of the *equilibrium* state.

1. \_\_\_\_\_
2. \_\_\_\_\_
3. \_\_\_\_\_
4. \_\_\_\_\_

**Chemistry 12****Unit 2 - Chemical Equilibrium**

5. Explain why the colour of NO<sub>2</sub> gas first gets **darker** and then gets *lighter* when compressed in a syringe. The equation is:




---

---

---

---

6. Define *enthalpy* \_\_\_\_\_
7. Define *entropy* \_\_\_\_\_
8. For the reaction:




The tendency toward *minimum enthalpy* favours the \_\_\_\_\_. The  
tendency toward *maximum entropy* favours the \_\_\_\_\_

If the reactants are combined will the reaction *go to completion, not occur at all* or *reach a state of equilibrium*?

---

9. For the reaction:



As this reaction proceeds right, enthalpy is \_\_\_\_\_creasing and entropy is  
\_\_\_\_\_creasing.

If the reactants are combined will the reaction *go to completion, not occur at all* or  
*reach a state of equilibrium*?

---

**Chemistry 12****Unit 2 - Chemical Equilibrium**

10. For the reaction:



The tendency toward *minimum enthalpy* favours the \_\_\_\_\_. The  
tendency toward *maximum entropy* favours the \_\_\_\_\_.

If the reactants are combined will the reaction *go to completion, not occur at all* or *reach a state of equilibrium*?

11. For the reaction:



Which way will the equilibrium shift when the following changes are made:

- a)  $\text{NaCl}_{(\text{aq})}$  is added ..... \_\_\_\_\_
- b) The pressure is increased ..... \_\_\_\_\_
- c) The  $[\text{OH}^-]$  is decreased ..... \_\_\_\_\_
- d) The temperature is decreased ..... \_\_\_\_\_
- e) The volume of the container is decreased ..... \_\_\_\_\_
- f) The solid sodium is chopped into smaller pieces.. \_\_\_\_\_
- g) A catalyst is added ..... \_\_\_\_\_

12. For the following reaction:




Which way will the equilibrium shift (if any) when the following changes are made:

- a)  $[\text{NO}]$  is decreased ..... \_\_\_\_\_
- b)  $[\text{O}_2]$  is increased ..... \_\_\_\_\_
- c)  $[\text{NH}_3]$  is increased ..... \_\_\_\_\_

**Chemistry 12****Unit 2 - Chemical Equilibrium**

- d) The temperature is decreased ..... \_\_\_\_\_
- e) The volume of the container is increased ..... \_\_\_\_\_
- f) The total pressure is increased ..... \_\_\_\_\_
- g) Helium gas is added to increase the total pressure ..... \_\_\_\_\_
- h) The temperature is increased ..... \_\_\_\_\_
- i) A catalyst is added ..... \_\_\_\_\_
13. Discuss the ideal pressure and temperature conditions for achieving maximum yield of ammonia at a reasonable rate in the *Haber Process*:




Pressure should be (*high/low*) \_\_\_\_\_ and temperature should be (*high/low*) \_\_\_\_\_, but still high enough to maintain a \_\_\_\_\_

Give reasons for your answer. \_\_\_\_\_

\_\_\_\_\_

14. In an experiment at 423°C, the following concentrations were measured for the ***equilibrium*** system:



$$[\text{HI}] = 17.7 \times 10^{-3} \text{ M}, \quad [\text{H}_2] = 1.83 \times 10^{-3} \text{ M} \quad \text{and} \quad [\text{I}_2] = 3.13 \times 10^{-3} \text{ M}.$$

Calculate the *value* for the equilibrium constant ( $K_{\text{eq}}$ ) at 423°C .


Answer \_\_\_\_\_

**Chemistry 12****Unit 2 - Chemical Equilibrium**

15. If, at 423°C , the  $[H_2]$  and  $[I_2] = 4.8 \times 10^{-3} \text{ M}$ , calculate the  $[HI]$ . Use  $K_{eq}$  from question 14.

Answer \_\_\_\_\_

16. Given the equilibrium equation:



If 2.0 moles of  $X_2$  and 2.0 moles of  $Y_2$  are added to a 1.0 L container, an equilibrium is established in which the  $[Y_2] = 0.80 \text{ M}$ . Find the following at equilibrium. (*Use a table.*)

$[X_2] =$  \_\_\_\_\_  $[XY_3] =$  \_\_\_\_\_  $K_{eq} =$  \_\_\_\_\_

**Chemistry 12****Unit 2 - Chemical Equilibrium**

17. The equation:  $A_{(g)} + B_{(g)} \rightleftharpoons C_{(g)} + D_{(g)}$  has a  $K_{eq} = 49$  at  $25^{\circ}\text{C}$ .

If 1.0 mole of C and 1.0 mole of D are added to a 0.50 L container at  $25^{\circ}\text{C}$ , calculate the following *at equilibrium*: (Use a table.)

[A] = \_\_\_\_\_ [B] = \_\_\_\_\_ [C] = \_\_\_\_\_ [D] = \_\_\_\_\_


18. In the equilibrium in question 17, what, if anything, would happen to the *value* of the equilibrium constant if the *temperature is increased*?

Answer \_\_\_\_\_

Explain your answer. \_\_\_\_\_

\_\_\_\_\_

19. Write the *K<sub>eq</sub> expression* for the following reaction: (*Be careful of phases!*)





**Chemistry 12****Unit 2 - Chemical Equilibrium**

20. The  $K_{eq}$  for the reaction:




Using this information, calculate the  $K_{eq}$  for the reaction :



Answer \_\_\_\_\_

21. Consider the following equilibrium:




What will occur when 1.0 mol of  $\text{SO}_2$ , 1.0 mol of  $\text{O}_2$ , and 1.0 mol of  $\text{SO}_3$  are placed in a 1.0 L container and allowed to reach equilibrium?

- a)  $[\text{SO}_2]$  will increase,  $[\text{SO}_3]$  will increase.
- b)  $[\text{SO}_2]$  will increase,  $[\text{SO}_3]$  will decrease.
- c)  $[\text{SO}_2]$  will decrease,  $[\text{SO}_3]$  will increase.
- d)  $[\text{SO}_2]$  will decrease,  $[\text{SO}_3]$  will decrease.

Answer \_\_\_\_\_

Explain your answer. \_\_\_\_\_  
\_\_\_\_\_

22. Given the equilibrium:



Give 5 actions which could cause this equilibrium to shift to the *left*:

- 1. \_\_\_\_\_
- 2. \_\_\_\_\_
- 3. \_\_\_\_\_
- 4. \_\_\_\_\_
- 5. \_\_\_\_\_



## Chemistry 12

## Unit 2 - Chemical Equilibrium

23. How does the addition of a *catalyst* affect the  $K_{eq}$  for a system? \_\_\_\_\_
24. Choose the equilibrium which most *favours the reactants*.
- |   |                                |
|---|--------------------------------|
| a) $\text{NO} + 1/2\text{O}_2 \rightleftharpoons \text{NO}_2$                   | $K_{eq} = 4.4 \times 10^7$     |
| b) $\text{CO} + 1/2\text{O}_2 \rightleftharpoons \text{CO}_2$                   | $K_{eq} = 4.0 \times 10^{-3}$  |
| c) $\text{C} + \text{H}_2\text{O} \rightleftharpoons \text{CO} + \text{H}_2$    | $K_{eq} = 3.1 \times 10^3$     |
| d) $\text{NO} + \text{H}_2\text{O} \rightleftharpoons \text{NO}_2 + \text{H}_2$ | $K_{eq} = 1.0 \times 10^{-22}$ |
25. Consider the following system:  $2\text{SO}_2(\text{g}) + \text{O}_2(\text{g}) \rightleftharpoons 2\text{SO}_3(\text{g})$   $\Delta H = -99 \text{ kJ/mol}$   
What are *four* things which could be done in order to increase the yield of  $\text{SO}_3$ ?
1. \_\_\_\_\_
  2. \_\_\_\_\_
  3. \_\_\_\_\_
  4. \_\_\_\_\_
26. Given the equilibrium equation:  $\text{XY}(\text{g}) + \text{heat} \rightleftharpoons \text{X}(\text{g}) + \text{Y}(\text{g})$

If initially, at equilibrium, the  $[\text{XY}] = 3.0 \text{ M}$ , the  $[\text{X}] = 5.0 \text{ M}$  and the  $[\text{Y}] = 6.0 \text{ M}$ , draw a graph showing qualitatively what happens to the concentrations of each species as the following changes are made to the system:

Time I - The *temperature is decreased*.

Time II - Some  $\text{X}(\text{g})$  is *removed from* the system

Time III - Some  $\text{XY}(\text{g})$  is *added to* the system

Time IV - The *total pressure is increased*.

